



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Solution by A. H. HOLMES, Brunswick, Me.

$$x^2 + w^2 + v^2 = a^2 \dots\dots\dots (1), \quad vw + u(y+z) = bc \dots\dots\dots (4),$$

$$w^2 + y^2 + u^2 = b^2 \dots\dots\dots (2), \quad wu + v(z+x) = ca \dots\dots\dots (5),$$

$$v^2 + u^2 + z^2 = a^2 \dots\dots\dots (3), \quad uv + w(x+y) = ab \dots\dots\dots (6).$$

From (2), (3), and (4),

$$(w^2 + y^2 + u^2)(v^2 + u^2 + z^2) = [vw + u(y+z)]^2.$$

$$\therefore (yv - uw)^2 + (uv - zw)^2 + (u^2 - yz)^2 = 0.$$

$$\therefore yv = uw \dots\dots\dots (7); \quad uv = zw \dots\dots\dots (8); \quad u^2 = yz \dots\dots\dots (9).$$

From (1), (3), and (5),

$$(x^2 + w^2 + v^2)(v^2 + u^2 + z^2) = [wu + v(z+x)]^2.$$

$$\therefore v^2 = xz \dots\dots\dots (10); \quad vw = xu \dots\dots\dots (11).$$

From (2), (3), and (6),

$$(w^2 + y^2 + u^2)(v^2 + u^2 + z^2) = [uv + w(x+y)]^2.$$

$$\therefore w^2 = xy \dots\dots\dots (12).$$

Substituting xy and xz for w^2 and v^2 in (1), xy and yz for w^2 and u^2 in (2), and xz and yz for v^2 and u^2 in (3), adding the three equations and extracting square root, we obtain $x+y+z = \sqrt{a^2 + b^2 + c^2}$.

$$\text{Substituting } xu \text{ for } vw \text{ in (4), } u = \frac{bc}{\sqrt{a^2 + b^2 + c^2}}.$$

$$\text{Similarly, } v = \frac{ac}{\sqrt{a^2 + b^2 + c^2}}, \quad w = \frac{ab}{\sqrt{a^2 + b^2 + c^2}},$$

$$x = \frac{a^2}{\sqrt{a^2 + b^2 + c^2}}, \quad y = \frac{b^2}{\sqrt{a^2 + b^2 + c^2}}, \quad z = \frac{c^2}{\sqrt{a^2 + b^2 + c^2}}.$$

227. Proposed by G. I. HOPKINS, A. M., Manchester, N. H.

$$\text{Solve } x + y + xy + x^2y + xy^2 + x^3y + 2x^2y^2 + xy^3 + x^3y^2 + x^2y^3 = 11; \quad x^4y + 3x^3y^2 + 3x^2y^3 + 2x^4y^2 + 4x^3y^3 + 2x^2y^4 + 4x^4y^3 + 4x^3y^4 + xy^4 + x^5y^2 + x^5y^3 + 2x^4y^4 + x^2y^5 + x^3y^5 = 30.$$

Solution by F. P. MATZ, Ph. D., Sc. D., Reading, Pa.

Put $X = (x + y + xy + x^2y + xy^2)$, and $Y = (x^3y + 2x^2y^2 + xy^3 + x^3y^2 + x^2y^3)$; then the given equations become, respectively,

$$X + Y = 11 \dots\dots\dots (\alpha), \text{ and}$$

$$XY = 30 \dots\dots\dots (\beta).$$

$$\therefore X = 6, \text{ or } 5; \text{ and } Y = 5, \text{ or } 6.$$

By putting $X_1=(x+y+xy)$, and $Y_1=(x^2y+xy^2)$, the expressions represented by X and Y give, respectively,

$$\begin{aligned} X_1+Y_1 &= 6, \text{ or } 5 \dots\dots\dots (\alpha_1), \text{ and} \\ X_1Y_1 &= 5, \text{ or } 6 \dots\dots\dots (\beta_1). \end{aligned}$$

$$\therefore X_1=5, 1, 3, \text{ or } 2; \text{ and } Y_1=1, 5, 2, \text{ or } 3.$$

By putting $X_2=(x+y)$ and $Y_2=(xy)$, the expressions represented by X_1 and Y_1 give, respectively,

$$\begin{aligned} X_2+Y_2 &= 5, 1, 3, \text{ or } 2 \dots\dots\dots (\alpha_2), \text{ and} \\ X_2Y_2 &= 1, 5, 2, \text{ or } 3 \dots\dots\dots (\beta_2). \end{aligned}$$

$$\begin{aligned} \therefore x+y &= \frac{1}{2}(5 \pm \sqrt{21}), \frac{1}{2}(1 \pm \sqrt{-19}), 2 \text{ or } 1, \text{ or } 1 \pm \sqrt{-2}; \text{ and} \\ xy &= \frac{1}{2}(5 \mp \sqrt{21}), \frac{1}{2}(1 \mp \sqrt{-19}), 1 \text{ or } 2, \text{ or } 1 \mp \sqrt{-2}. \end{aligned}$$

Solving these *eight* simultaneous equations, we have the *sixteen* values of x and y .

Also solved by J. Scheffer, Henry Heaton, G. B. M. Zerr, and A. H. Holmes.

GEOMETRY.

254. Proposed by W. J. GREENSTREET, M. A., Editor of the Mathematical Gazette, Stroud, England.

Find the cartesian equation to a line that is both tangent and normal to the cardioid.

[No solution of this problem has been received.]

255. Proposed by J. SCHEFFER, A. M., Hagerstown, Md.

Find the envelope of the straight line that connects the extremities of two conjugate diameters of an ellipse.

I. Solution by G. W. GREENWOOD, M. A., Professor of Mathematics, McKendree College, Lebanon, Ill.

If we project the ellipse into a circle, the conjugate diameters are projected into perpendicular diameters of the circle, whose chord envelopes a concentric circle. Hence in the original figure the chord envelopes a similar, and similarly situated concentric ellipse.

II. Solution by W. J. GREENSTREET, M. A., Editor of The Mathematical Gazette, Stroud, England.

The equation to the line is, with usual notation,

$$\frac{x}{a} \cos(\varphi + \frac{1}{4}\pi) + \frac{y}{b} \sin(\varphi + \frac{1}{4}\pi) = \cos \frac{1}{4}\pi, \text{ or } \left(\frac{x}{a} + \frac{y}{b}\right) \cos \varphi - \left(\frac{x}{a} - \frac{y}{b}\right) \sin \varphi = 1,$$

$$i. e., \left(\frac{x}{a} + \frac{y}{b} - 1\right) - 2\left(\frac{x}{a} - \frac{y}{b}\right) \tan \frac{1}{2}\varphi - \left(\frac{x}{a} + \frac{y}{b} + 1\right) \tan^2 \frac{1}{2}\varphi = 0.$$

\therefore The equation to the envelope is